


25/30

الرقم الجامعي: _____ الاسم: _____
رقم الشعبة أو وقت المحاضرة: _____ اسم الدكتور: _____
Consider $g = 9.8 \text{ m/s}^2$

Question	1	2	3	4	5	6	7	8	9	10	11	12
Answer	a B	b	c	d	e	e	d	b	a	c	A	e

- ✓ 1. An applied force of 10 N compresses (يضغط) a spring with a 20-N/m spring constant. The work done (in J) by this force is:
 (A) 0.5 B) 2.5 C) 5 D) 10 E) 200


- ✓ 2. Three blocks (A, B, C), each having the same mass M , are connected by strings as shown. Block C is pulled to the right by a force \vec{F} that causes the entire system to accelerate. Neglecting friction, the net force acting on block B is:



- A) 0 (B) $\vec{F}/3$ (C) $\vec{F}/2$ D) $2\vec{F}/3$ E) \vec{F}
- ✓ 3. A constant force (50-N) acts on a 2-kg box that starts from rest. When the force has been acting for 2 s the rate at which it is doing work (in W) is:
 A) 100 B) 1000 (C) 2500 D) 5000 E) 63000

- ✓ 4. Let M denote the mass of Earth and let R denote its radius. The ratio g/G at Earth's surface is:
 A) R^2/M B) M/R C) MR^2 (D) M/R^2 E) R/M

- ✓ 5. The ball shown is being swung (يتأرجح) in a vertical circle at the end of a 0.7-m string. The minimum speed (in m/s) needed for the ball to pass successfully over the top position is:




- A) 1.3 B) 9.8 C) 3.9 D) 6.9 (E) 2.6
- ✓ 6. A 40-N box rests on a rough (خشن) horizontal floor. A 12-N horizontal force is then applied to it. If the coefficients of friction are $\mu_s = 0.5$ and $\mu_k = 0.4$, the magnitude of the frictional force on the box (in N) is:
 A) 8 B) 40 C) 16 D) 20 (E) 12

- ✓ 7. The energy transferred to a system as a function of time is equal to:
 $[E(t) = 3.5 t + 6.2 t^2, \text{ (in J)}]$. The time rate of this energy (in W) at $t = 3.1 \text{ s}$ is:
 A) 3.5 B) 6.2 C) 16 (D) 42 (E) 70

- ✓ 8. The system shown remains at rest.
The force of friction (in N) on the block on the incline is:


A) 4 **B) 8** C) 12
D) 16 E) 20



9. As a 2.0-kg object moves from $(2\hat{i} + 5\hat{j})$ m to $(6\hat{i} - 2\hat{j})$ m, the constant resultant force acting on it is equal to $(4\hat{i} - 3\hat{j})$ N. If the speed of the object at the initial position is 4.0 m/s, its kinetic energy (in J) at the final position is:


A) 53 B) 62 C) 73 D) 86 E) 24

- ✓ 10. The plot below shows the force on an object as it moves along the x axis.
The work (in J) done on the object as it moves from $x = 0$ m to $x = 20$ m is:



A) 40 B) 90 **C) 450** D) 200 E) 750

- ✓ 11. A box with a weight of 50 N rests on a horizontal surface. A person pulls horizontally on it with a force of 15 N and it does not move. To start it moving, a second person pulls vertically upward on the box. If the coefficient of static friction is 0.4, the smallest vertical force (in N) for which the box moves is:



A) 12.5 **B) 5.5** C) 20.5 D) 25.5 E) 35.5

- ✓ 12. A 0.50-kg object moves on a horizontal frictionless circular track with a radius of 2.5 m. An external constant force of 3.0 N, always tangent to the track, causes the object to speed up as it goes around. If the object starts from rest, then at the end of one revolution (دورة) the radial component of the force (in N) of the track on it is:

A) 19 B) 96 C) 47 D) 75 **E) 38**

القوة المركزية
في دورتها

22
30

اسم مدرس المادة:
وقت المحاضرة:
الرقم العتململ:

The University of Jordan
Faculty of Science
Physics Department

General Physics (b0302101)
Second Exam
Second Semester 2017/2018

Name (in Arabic): _____ Student ID: _____ Section: _____

(Exam duration: 75 minutes)

Note 1: Following are 15 multiple-choice questions. Write the symbol of correct answer in the answers table. Only the answers in the table will be graded.

Note 2: Ignore air resistance in all problems and take $|g| = 9.8 \text{ m/s}^2$ at the Earth's surface.


Answers table

Question number	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Symbol of correct answer	d	e	e	b	a	b	b	d	d	a	c	g	a	e	b


Q.1: The only three forces that act on a 3-kg particle are as follows: $\vec{F}_1 = (2\hat{i} + 3\hat{j})N$, $\vec{F}_2 = (\hat{i} + 2\hat{j})N$ and $\vec{F}_3 = (2\hat{i} + 5\hat{k})N$. The magnitude (in m/s^2) of the particle's acceleration is:
 a. 9.80 b. 4.33 c. 12.12 **d. 2.89** e. 20.46

Q.2: A force $\vec{F} = (6\hat{i} - 2\hat{j})N$ acts on a particle that undergoes a displacement $\Delta\vec{r} = (3\hat{i} - \hat{j})m$. The work (in Joules) done by this force on the particle is:
 a. 11 b. 14 c. 16 d. 18 **e. 20**

Q.3: A 2-kg hanging mass (m_1) is connected by a string over a pulley to a 20-kg block (m_2) that is sliding on a 50° fixed inclined plane (see the adjacent figure). If the pulley's mass and the mass of the string are negligible, and all surfaces are frictionless, the magnitude of the acceleration (in m/s^2) of the moving system is:
 a. 2.56 b. 9.80 c. 0.21
 d. 1.15 **e. 5.93**




Q.4: Two blocks $M_1 = 3 \text{ kg}$ and $M_2 = 5 \text{ kg}$ are in contact with each other on a frictionless, horizontal surface, as shown in the adjacent figure. If a horizontal force $F = 16 \text{ N}$ is applied to M_1 , the magnitude (in N) of the contact force between the two blocks is:
 a. 2 **b. 4** c. 7 d. 10 e. Zero



Q.5: An object of mass m , speed V and initial kinetic energy K_i . If the speed of the object becomes $3V$, then the ratio (K_f/K_i) is:
a. 9 b. (1/9) c. 1 d. 18 e. 81

Q.6: A 50-kg object slides from rest from point A on the rough track shown in the adjacent figure. If the speed of the particle at point B is 6 m/s. The work (in J) done by frictional forces is:

- a. -7300 b. -8900 c. -3700
d. -4000 e. -5300



Q.7: A ball of mass 2 kg is fired straight up with an initial speed of 20 m/s. It rises to its maximum height, and then falls down to its starting point. Neglecting air resistance, the work (in J) done on the ball by gravitational force through the entire trip is:

- a. -22.8 b. 18.6 c. zero d. 22.8 e. -18.6


Q.8: The adjacent figure shows a setup of three masses that are connected by three wires. The whole system is under static equilibrium. If $m_1 = 15$ kg, $m_2 = 25$ kg and $m_3 = 60$ kg. The tension (T_1) in the first wire (measured in Newtons) is:

- a. 680 b. 588 c. 196 d. 980 e. 294



Q.9: The adjacent figure shows a box of mass 3 kg moving on a horizontal, frictionless surface with a speed of 4 m/s towards an unstretched spring of negligible mass that is attached horizontally to a rigid wall. The box collides with the spring and stops momentarily before reversing direction. If the spring constant is 1000 N/m, the maximum compression (in m) of the spring is:

- a. 0.550 b. 0.219 c. 0.357
 d. 0.179 e. 0.742



Q.10: True or False:

"The work done by any conservative force on a particle moving through any closed path is zero"

- a. True b. False

Q.11: A box with initial speed $V_i = 5$ m/s slides on a rough horizontal surface. If the coefficient of kinetic friction is 0.8, the distance (in m) moved by the box before coming to a stop is:

- a. 0.56 b. 2.34 c. 1.59 d. 3.14 e. 8.43

Q.12: A potential energy function for a two-dimensional force is of the form:


$U(x, y) = (3x^2y - 7x)$ J. The magnitude of the force (in N) that acts at the point (1, 2) m is:

- a. 5.83 b. 3.77 c. 9.80 d. 12.65 e. 25.41

$6xy - 7$
 $5j$
 $3i^2$
 $3i - 3$


Q.13: A block of mass M rests on an inclined rough surface. The inclination angle of the surface is increased to θ_c at which point the block becomes on the verge of slipping. The coefficient of static friction of the surface is:

- a. $\sin(\theta_c)$ b. $\cos(2\theta_c)$ c. $\sin(4\theta_c)$
 d. $\tan(\theta_c)$ e. $\sin^2(\theta_c)$



Q.14: A small ball of mass m is suspended from a string of length L . The ball revolves with constant speed v in the horizontal circle of radius r as shown in the adjacent figure. If the string makes an angle θ with the vertical direction, the speed v of the ball is given by:

- a. $\sqrt{rg \sin \theta}$ b. $\sqrt{rg \cos \theta}$
 c. $\sqrt{rg \csc \theta}$ d. $\sqrt{rg \cot \theta}$
 e. $\sqrt{rg \tan \theta}$



Q.15: A force $\vec{F} = (8\hat{i} + 3\hat{j})$ N acts on a box that is sliding on a floor. At the instant the velocity of the box is $\vec{V} = (3\hat{i} - 2\hat{j})$ m/s, the instantaneous power (in Watts) supplied by this force is:


- a. 30 b. 18 c. 11 d. 55 e. 78

Good Luck!!!

Question	1	2	3	4	5	6	7	8	9	10	11	12
Answer	D	F	A	B	C	D	F	F	D	A	B	A

1. An applied force of 10 N compresses (يضغط) a spring with a 20-N/m spring constant. The work done (in J) by this force is:
 A) 5 B) 200 C) 0.5 **D) 10** E) 2.5

2. Three blocks (A, B, C), each having the same mass M , are connected by strings as shown. Block C is pulled to the right by a force \vec{F} that causes the entire system to accelerate. Neglecting friction, the net force acting on block B is:



A) 0 B) \vec{F} C) $\vec{F}/2$ D) $2\vec{F}/3$ **E) $\vec{F}/3$**

3. A constant force (50-N) acts on a 2-kg box that starts from rest. When the force has been acting for 2 s the rate at which it is doing work (in W) is:
A) 2500 B) 1000 C) 100 D) 5000 E) 63000

4. Let M denote the mass of Earth and let R denote its radius. The ratio g/G at Earth's surface is:
 A) R^2/M **B) M/R^2** C) MR^2 D) M/R E) R/M

5. The ball shown is being swung (بفارجح) in a vertical circle at the end of a 0.7-m string. The minimum speed (in m/s) needed for the ball to pass successfully over the top position is:



A) 1.3 B) 9.8 **C) 2.6** D) 6.9 E) 3.9

6. A 40-N box rests on a rough (خشن) horizontal floor. A 12-N horizontal force is then applied to it. If the coefficients of friction are $\mu_s = 0.5$ and $\mu_k = 0.4$, the magnitude of the frictional force on the box (in N) is:

A) 8 B) 40 C) 12 **D) 20** E) 16

7. The energy transferred to a system as a function of time is equal to:
 $[E(t) = 3.5t + 6.2t^2, \text{ (in J)}]$. The time rate of this energy (in W) at $t = 3.1$ s is:
 A) 3.5 B) 42 C) 16 D) 6.2 **E) 70**

22
30

اسم مدرس المادة:
رقم الشعبة: ٢٠٠
الرقم المتسلسل: 21

The University of Jordan
Faculty of Science
Physics Department

General Physics (1) (0302101)
Second Exam
First Semester 2016/2017

• Student's Name: ...
Student's ID: ...

Note 1: Following are 10 multiple-choice questions. Write the symbol of correct answer in the answers' table. **Only** the answers in the table will be graded.

Note 2: Ignore air resistance in all problems and take $|g| = 9.8 \text{ m/s}^2$ at the Earth's surface.

Note 3: The significant digit notation is not taken into account throughout the given answers.

Answers' Table

Question Number	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Symbol of Correct Answer	b	b	c	e	b	e	d	b	a	b	e	c	c	a	a

Q.1: A force $\vec{F} = (2\hat{i} - \hat{j})N$ acts on an object. The work (in J) that this force does as the object moves from the origin (0, 0, 0) to the point (13, 11, 0) m is:

[Hint: the displacement of the particle is: $\Delta \vec{r} = (13\hat{i} + 11\hat{j})m$]

- a. 246 **b. 26** c. 37 d. 15 e. 100

Q.2: The work performed as a function of time for a process is given by $W = at^3$, where $a = 2.4 \text{ J/s}^3$. The instantaneous power output (measured in W) at $t = 8.7 \text{ sec}$ is:

- a. 138 **b. 545** c. 125 d. 207 **e. 912**

Q.3: In the adjacent figure, two identical ideal massless springs have unstretched lengths of 0.25 m and spring constants of 550 N/m. The springs are attached to a small cube and stretched to a length L of 0.30 m as in Figure A. An external force P pulls the cube a distance $D = 0.020 \text{ m}$ to the right and holds it there. (See Figure B.) The external force P , that holds the cube in place in Figure B, is:

- a. 34 N b. 45 N **c. 28 N**
d. 22 N e. 11 N

Figure A





Figure B




Q.4: A force $F = bx^3$ acts in the x direction, where the value of b is 3.7 N/m^3 . The work (in J) done by this force in moving an object from $x = 0.00 \text{ m}$ to $x = 2.6 \text{ m}$ is:

- a. 98.4 b. 27.3 c. 50.4 d. 9.8 **e. 42.2**


Q.5: In the adjacent figure, a block of mass M hangs at rest. The rope that is fastened to the wall is horizontal and has a tension of 52 N. The rope that is fastened to the ceiling has a tension of 104 N, and makes an angle θ with the ceiling. The angle θ (measured in degrees) is:

- a. 55° b. 60° c. 30° d. 85° e. 15°



Q.6: A weight $W_1 = 20$ N rests on a second weight $W_2 = 50$ N on a perfectly smooth horizontal floor as shown in the adjacent figure. When a horizontal force $F = 15$ N is applied on the lower box (see adjacent figure), both boxes move together. The magnitude (in N) and direction of the net external force on the upper box is:

- a. 6.48 N to the right b. 6.48 N to the left
c. 4.28 N to the left d. 4.28 N to the right
e. Zero




Q.7: A 5.00-kg box slides 9.00 m across a horizontal floor before coming to rest. If the box had an initial speed of 3.00 m/s, then, the coefficient of kinetic friction (μ_k) between the floor and the box is:

- a. 0.412 b. 0.587 c. 0.321 d. 0.051 e. 0.115


Q.8: A system comprising blocks, a light frictionless pulley, a frictionless incline, and connecting ropes is shown in the adjacent figure. The 9.0-kg block accelerates downward when the system is released from rest. The tension in the rope connecting the 6.0-kg block and the 4.0-kg block (measured in N) is:

- a. 80 b. 12 c. 42 d. 99 e. 60



Q.9: Two objects are connected by a very light flexible string that passes over a very light and frictionless pulley as shown in the adjacent figure. Neglecting air resistance. If $M = 0.60$ kg and $m = 0.40$ kg, the tension in the string (measured in N) is:

- a. 4.7 b. 21.1 c. 14.3 d. 9.8 e. 19



Q.10: True or False:

“The action and reaction forces are equal in magnitude, opposite in direction and act on the same objects”

True

False

Q.11: Two moons orbit a planet in nearly circular orbits. Moon A has orbital radius r , and moon B has orbital radius $4r$. Moon A takes 15 days to complete one orbit. Neglecting gravitational interactions between the two moons, the time (measured in days) needed for moon B to complete an orbit is:

- a. 360 b. 180 c. 80 d. 160 e. 120

Q.12: Planet X has a mass equal to $1/3$ that of Earth, a radius equal to $1/3$ that of Earth, and an axial spin rate $1/2$ that of Earth. With g representing, as usual, the acceleration due to gravity on the surface of Earth, the acceleration due to gravity on the surface of planet X is:


- a. $g/3$ b. $g/9$ c. $3g$ d. $6g$ e. $9g$

Q.13: A block is on a frictionless horizontal table, on earth. This block accelerates at 3 m/s^2 when a 90 N horizontal force is applied to it. The block and table are then set up on the moon where the acceleration due to gravity is 1.62 m/s^2 . The weight (measured in N) of the block on the moon is:

- a. 93.7 b. 76.7 c. 48.6 d. 28.2 e. 36.8

Q.14: The adjacent figure shows a setup of three masses that are connected by three wires. The whole system is under static equilibrium. If $m_1 = 10 \text{ kg}$, $m_2 = 20 \text{ kg}$ and $m_3 = 70 \text{ kg}$, The tension (T_1) in the first wire (measured in Newtons) is:

- a. 980 b. 518 c. 294 d. 426 e. 686



Q. 15: True or False:

“Any non-accelerating frame of reference is considered as an inertial reference frame”

- a. True b. False

Good Luck!!!

Lecturer's Name:

أ.د. أم نعمة

Time: 16:00–17:10

*Take $g = 9.8 \text{ m/s}^2$ and $G = 6.7 \times 10^{-11} \text{ N.m}^2/\text{kg}^2$.

** Fill in the Table at the END with your answers, using CAPITAL letters ONLY.

Q1) Only two forces act on a 5.0-kg mass. These are $F_1 = (2i - 4j) \text{ N}$ and $F_2 = (3i - 6j) \text{ N}$. The magnitude of the resulting acceleration (in m/s^2) is:

- (A) 1.0 (B) 2.0 (C) 5.0 (D) 0.22 (E) 2.2

Q2) A 5.0-kg mass is suspended (علقت) by a string from the ceiling (سقف) of an elevator. The tension in the string is 50 N. The acceleration (in m/s^2) of the elevator is:

- (A) 9.8, downward (B) 9.8, upward (C) 0.20, upward (D) 2.0, upward (E) 2.0, downward

Q3)



In the above figure, the surfaces are frictionless and force $P = 10 \text{ N}$. The magnitude of the force (in N) exerted (المؤثرة) on block 1 by block 2 is:

- (A) 10 (B) 8.0 (C) 6.0 (D) 4.0 (E) 2.0

Q4) A block is released from rest on a 30° -incline and slides 9.0 m in 3.0 s. What is the coefficient of kinetic friction between the block and the surface of the incline?

- (A) 0.17 (B) 0.81 (C) 0.34 (D) 0.28 (E) 0.22

Q5) A mass of 1.0 kg, attached to the end of a string, swings in a vertical circle of radius 2.0 m. When the mass is at the lowest point of the circle, its speed is 10 m/s. The tension (in N) in the string at this point is:

- (A) 60 (B) 40 (C) 30 (D) 20 (E) 10

Q6) A point is at a distance $4R_E$ above the surface of the Earth (R_E being the Earth's radius which you need *not* know). The magnitude of the free-fall acceleration (in m/s^2) at this point is:


- (A) 9.8 (B) 2.0 (C) 2.5 (D) 0.39 (E) 0.61

Q7) The initial velocity of a 5.0-kg particle is $(2.0i - 5.0j)$ m/s. After t s, the velocity becomes $((5.0i - 6.0j)$ m/s. The work done (in J) by the *resultant* force during this time interval is:

- (A) zero (B) 100 (C) 80 (D) 10 (E) 425

Q8) A particle moves along the x-axis. It is acted upon by a force F_x (in N) that varies with position x (in m) as shown in the graph below. What work (in J) is done by this force as the particle moves from $x = 2$ m to $x = 12$ m?


- (A) +40 (B) -30 (C) +30 (D) -10 (E) +10



Q9) A 1.0-kg block slides (يتزلق) down a 30° -incline at a constant speed of 10 m/s. At what rate (in W) is work done on the block by the gravitational force?

- (A) +49 (B) -98 (C) zero (D) +100 (E) -100

Q10)



In the above figure, the system is released from rest with the spring in its equilibrium position. The pulley and the horizontal surface are frictionless. If the spring constant $k = 600 \text{ N/m}$ and $M = 5.0 \text{ kg}$, what is the maximum extension (امتداد) (in cm) of the spring?

- (A) 50 (B) 16 (C) 80 (D) 24 (E) 20

Q11) A 5.0-kg particle is dropped from rest. After falling a distance of 100 m, it has a speed of 25 m/s. What is the work done (in kJ) by the nonconservative air-resistive force on the particle during this fall?

- (A) +2.0 (B) -2.5 (C) +2.9 (D) -3.3 (E) -3.9

Q12) The potential energy function for a certain system is given (in J) by the expression $U(x,y) = x^2y^4 - 4x + 3y$, where x and y are in m. The magnitude of the corresponding force (in N) at $x = 1.0 \text{ m}$ and $y = 1.0 \text{ m}$ is:

- (A) zero (B) 6.0 (C) 9.0 (D) 7.0 (E) 7.3

Fill in the Table below with your answers, using CAPITAL letters ONLY:

Q1	Q2	Q3	Q4	Q5	Q6	Q7	Q8	Q9	Q10
E	C	E	A	A	E	C	C	A	A
				Q11	Q12				
				D	E				



THE UNIVERSITY OF JORDAN

PHYSICS DEPARTMENT

GENERAL PHYSICS I (0302101) / SECOND EXAM / AUG 6th 2016

SUMMER SEMESTER 2015/2016

 25
 /
 30

الرقم الجامعي:

رقم الشعبة:

اسم الطالب:

اسم العنبر:

	A	B	C	D	E		A	B	C	D	E		A	B	C	D	E
Q1	✓	✓				Q5	✓	✓				Q9		✓			
Q2		✓	✓			Q6	✓	✓				Q10					✓
Q3		✓	✓			Q7				✓		Q11		✓	✓		
Q4	✓	✓				Q8	✓	✓				Q12		✓	✓		


ANSWER ALL THE FOLLOWING QUESTIONS

- A mass of 25 kg is acted on by two forces: force F_1 is 25 N due east, and force F_2 is 15 N due north. The magnitude of the acceleration (in m/s^2) of the mass is:
 a) 1.17 b) 2.41 c) 3.63 d) 2.15 e) 4.26
- A 0.20-kg object attached to the end of a string swings in a vertical circle (radius = 80 cm). At the top of the circle the speed of the object is 4.5 m/s. The magnitude of the tension (in N) in the string at this position is:
 a) 2.0 b) 3.1 c) 6.4 d) 7.6 e) 18.8
- A force accelerates a body of mass M . The same force applied to a second body produces three times the acceleration. The mass of the second body is:
 a) M b) $3M$ c) $M/3$ d) $9M$ e) $M/9$
- It takes 32.0 J of work to stretch a spring 20.0 cm from its unstressed length. The extra work (in J) required to stretch the spring an additional 10.0 cm is:
 a) 16 b) 14 c) 40 d) 12 e) 51
- The required work (in kJ) for a 2000 kg car moving on a horizontal road to increase its velocity from $(2i + 3j)$ m/s to $(5i + 12j)$ m/s is:
 a) 156 b) 182 c) 312 d) 243 e) 426.3
- A constant resultant force of $(2i + 5j)$ N affects an object and causes it to move with constant velocity of $4j$ m/s. The power (in Watts) required for this process is:
 a) 20 b) 12 c) 42 d) 28 e) 17



7. A 25.0 kg block is initially at rest on a horizontal surface. A horizontal force of 75.0 N is required to set the block in motion. After it is in motion, a horizontal force of 60.0 N is required to keep the block moving with constant speed. The coefficient of kinetic friction is:
 a) 0.31 b) 1.0 c) Zero **d) 0.24** e) 0.1
8. A force acting on an object moving along the x axis is given by $F = (14x - 3.0x^2)$ N, where x is in m. The work (in Joules) done by this force as the object moves from $x = -1$ m to $x = 2$ m is:
a) 12 b) 28 c) 40 d) 42 e) -28
9. A 2.0 kg mass is projected from the edge of the top of a 20 m tall building with a velocity of 24 m/s. Ignoring air resistance, the kinetic energy (in kilo Joules) of the mass just before it strikes the leveled ground is:
 a) 0.18 **b) 0.97** c) 0.89 d) 0.26 e) 0.4
10. A 700 N university student in basic training climbs a 10.0 m vertical rope at a constant speed in 8.00 s. His power output (in Watts) is:
 a) 560 b) 600 c) 900 d) 700 **e) 875**
11. A potential energy function for a two-dimensional force is of the form $U = 3x^2y - 7x$. The force components that act at the point (1, 0) are:
 a) (-7, 2) b) (3, -7) **c) (7, -3)** d) (-3, 7) e) (0, 0)
12. The figure represents the total acceleration of a particle moving clockwise in a circle of radius 2.50 m at a certain time. At this instant, the tangential acceleration (in m/s^2) is:

- a) 13
 b) 5.7
c) 7.5
 d) Zero
 e) 15



The University of Jordan / Department of Physics
First Semester 2015/2016
Physics 101/ Second Exam

Section number : KEY

Student name (بالعربية): _____

Lecturer name : _____

Student number : _____

✓ Some helpful information: gravitational acceleration $g = 9.8 \text{ m/s}^2$

Notes: Turn off your cell phone and put it out of sight. Keep your calculator on your own desk. Calculators cannot be shared. You have 75 minutes to complete your exam. Be sure to fill the box below with your final answers before the end of the exam.

	A	B	C	D	E		A	B	C	D	E
1						7					
2						8					
3						9					
4						10					
5						11					
6						12					

- A particle of mass (11 kg) is subject to two forces such that one force has a magnitude of 21 N directed east, and the other force has a magnitude of 39 N directed east-north, what is the magnitude of the particle's acceleration (in m/s^2)?
 (A) 2.8 **(B) 5.1** (C) 7.5 (D) 3.7 (E) 12
- An object of mass 4.0-kg is placed on top of an elevator floor. If the force exerted by the floor on the object is equal to 38 N. What is the acceleration of the elevator (in m/s^2)?
 (A) 0.8 upward (B) 0.8 downward (C) 1.3 upward (D) 1.3 downward **(E) 0.3 downward**
- A force of magnitude 20N directed in the positive x direction is acting on a particle and displacing it from the point (2m, -1m) to the point (4m, -3m). What is the work done by the force (in J)?
 (A) 60 **(B) 40** (C) 30 (D) 80 (E) 70
- A certain pendulum consists of a 1.5-kg mass swinging at the end of a string (length = 2.0 m). At the lowest point in the swing the tension in the string is equal to 20 N. To what maximum height (in cm) above this lowest point will the mass rise during its oscillation?
(A) 36 (B) 20 (C) 30 (D) 28 (E) 17
- A spring ($k = 600 \text{ N/m}$) is placed in a vertical position with its lower end supported by a horizontal surface. The upper end is compressed 20 cm, and a 4.0 kg block is placed on the compressed (مضغوط) spring. The system is then released from rest. How far above the point of release will the block rise (in cm)?
 (A) 20 **(B) 31** (C) 10 (D) 15 (E) 25

6. A potential energy function for a two-dimensional force is of the form $U = 3x^2y$. Find the force that acts at the point (1, 1).

- (A) $\vec{F} = -12\hat{i} - 3\hat{j}$ (B) $\vec{F} = -6\hat{j}$ (C) $\vec{F} = -24\hat{i} - 12\hat{j}$ (D) $\vec{F} = -6\hat{i} - 3\hat{j}$ (E) $\vec{F} = -6\hat{i}$

7. A 6.0-kg block slides along a horizontal surface. If $\mu_k = 0.20$ for the block and surface, at what rate is the friction force doing work on the block (in W) at an instant when its speed is 4.0 m/s?

- (A) -63 (B) -47 (C) +50 (D) +25 (E) -55


8. A particle of mass (1.5 kg) is moving on the x-axis with an acceleration given as $a = (6.0x + 5.0) \text{ m/s}^2$, What is the speed of the particle in (m/s) at the moment it reaches $x = 4.0 \text{ m}$, given that the particle started motion from origin with initial velocity 2.0 m/s?

- (A) 10.1 (B) 14.7 (C) 11.8 (D) 13.1 (E) 9.5

9. An airplane moves at constant speed of 140 m/s as it travels around a vertical circular loop which has a 1.0-km radius. What is the magnitude of the net force causing the centripetal acceleration on the 71-kg pilot (in N)?


- (A) 1000 (B) 1392 (C) 1200 (D) 1310 (E) 1022

10. A roller-coaster car has a mass of 400 kg when fully loaded with passengers (ركاب). At the bottom of a circular dip of radius 40 m (as shown in the figure) the car has a speed of 16 m/s. What is the magnitude of the force the track exerts on the car at the bottom of the dip (in kN)?



- (A) 10.1 (B) 9.7 (C) 8.1 (D) 13.1 (E) 6.5

11. What is the magnitude of the tension in the string (in N) if $M = 2.0 \text{ kg}$ in the figure shown? Assume the surface is frictionless.



- (A) 21. (B) 19.7 (C) 32.2 (D) 42.9 (E) 56.5

12. A box of mass (42 kg) is placed on top of a rough horizontal surface whose coefficients of friction are ($\mu_s = 0.6, \mu_k = 0.4$). If a man tried to push the box by applying a force of (210 N), what would be the magnitude of the friction force (in N)?

- (A) 210 (B) 247 (C) 220 (D) 165 (E) 230

Student's Name:

Student's Number:

Lecturer's Name:

Time: One Hour


Take the value of the gravitational acceleration $g = 9.8 \text{ m/s}^2$.

Fill in the Table (page 2) with the letters corresponding to your answers.

1. A particle is moving in a circular path. If F_r is the magnitude of the net force in the radial direction and F_t is the magnitude of the net tangential force acting on the particle then the total net force is


- A) $F = \sqrt{F_r + F_t}$ B) $F = F_r - F_t$
 C) $F = \sqrt{F_r^2 - F_t^2}$ D) $F = \sqrt{F_r^2 + F_t^2}$
 E) None of the above.

2. A small object of mass m starts at rest at the position shown and slides along the frictionless loop-the-loop track of radius R . What is the smallest value of y such that the object will slide without losing contact with the track?



- A) $R/2$ B) $R/4$ C) R D) $2R$ E) 0

3. A man moves the 15.0 g object shown in a vertical plane at a constant speed from position X to position Y along a circular track of radius 20.0 m. The process takes 0.75 min. The work done by the man is about:



- A) 0.0 J B) 5.9 J C) 11.8 J D) 2.0 J E) 3.9 J


4. A 0.4 kg particle moves along the x -axis under the influence of a conservative force. The potential energy is given by $U(x) = 8.0(\text{J/m}^2)x^2 + 4.0(\text{J/m}^4)x^4$ (J), where x is in meters. If the particle has a speed of 5.0 m/s when it is at $x = 1.0$ m, its speed when it is at the origin is:

- A) 2.5 m/s B) 5.6 m/s C) 9.2 m/s D) 11.2 m/s E) 8.7 m/s

5. The potential energy of a body of mass m is given by $U(x) = mgx + \frac{1}{2}kx^2$, where k is a constant. The magnitude of the corresponding force is:

- A) $\frac{1}{2}mgx^2 - \frac{1}{6}kx^3$ B) $-mg + \frac{1}{2}kx$ C) $mg - kx$
 D) $-\frac{1}{2}mgx^2 + \frac{1}{6}kx^3$ E) $-mg - kx$

6. An escalator is used to move 10 people (60 kg each) per minute from the first floor of a store to the second floor, 5 m above. The power required is approximately:




- A) 490 W B) 1 960 W C) 58 800 W D) 196 W E) 980 W

7. A nonconservative force (Hint think of the force of friction):
- A) cannot do any work
 - B) must be perpendicular to the velocity of the particle on which it acts.
 - C) violates Newton's third law.
 - D) violates Newton's second law.
 - E) none of the above.

8. A 6.0 kg block is released from rest 80 m above the ground. When it has fallen 50 m its kinetic energy is approximately (ignore air resistance):
- A) 1 176 J B) 47 040 J C) 2 940 J D) 196 J E) 3 528 J

9. The momentum of a system of particles is changing with time as $0.7t + 1.2t^2$, in kg m/s where t is in seconds. The magnitude of the net force at $t = 3.0$ s
- A) cannot be determined without knowing the momentum at $t = 0$.
 - B) is 7.9 N
 - C) is 3.1 N
 - D) is 5.5 N
 - E) cannot be determined without knowing the masses of the particles.

10. The figure shows a 3.00 kg steel ball which strikes a wall with a speed of 10.0 m/s at an angle of $\theta = 60^\circ$ with the surface. The ball bounces off with the same speed and angle. The ball is in contact with the wall for 0.20 s. The impulse is



- A) $\vec{I} = 260.0 \hat{j}$ (kg. m s⁻¹)
- B) $\vec{I} = -260.0 \hat{i}$ (kg. m s⁻¹)
- C) $\vec{I} = 52.0 \hat{i}$ (kg. m s⁻¹)
- D) $\vec{I} = -52.0 \hat{i}$ (kg. m s⁻¹)
- E) $\vec{I} = -52.0 \hat{j}$ (kg. m s⁻¹)

11. An object of mass $m_1 = 4.0$ kg is traveling at 6.0 m/s. It strikes an object of mass $m_2 = 8.0$ kg, which is stationary. The two objects stick together. Their common final speed is:

- A) 1.0 m/s B) 2.3 m/s C) 2.0 m/s D) 1.5 m/s E) 3.0 m/s

12. An object of mass $m_1 = 4.0$ kg is traveling at 3.0 m/s. It strikes an object of mass $m_2 = 8.0$ kg, which is stationary, in a head-on (elastic) collision. If v_{1f} and v_{2f} are the final velocities of m_1 and m_2 , after the collision, respectively, then $\left| \frac{v_{2f}}{v_{1f}} \right|$ is:

- A) 2.0 B) 1.5 C) 1.0 D) 0.5 E) 0.67

Question	1	2	3	4	5	6	7	8	9	10	11	12
Answer												

Q1 :



$$F_{net}^2 = F_c^2 + F_g^2$$

$$F_{net} = \sqrt{F_c^2 + F_g^2}$$

D

Q2 :

$$\Delta U + \Delta K = 0$$

$$-mgy + \frac{1}{2}m(v_f^2 - v_i^2) = 0$$

$$y = \frac{v_f^2}{2g}$$

$$= \frac{rg}{2g} = \frac{r}{2} \quad \underline{A}$$



$$N + mg = \frac{mv^2}{r}$$

$$N = 0 \Rightarrow v^2 = rg$$

Q3 :

$$W = -\Delta U$$

$$\Delta U = mgh$$

$$= 15 \times 40 \times 9.8 \times \frac{10^{-3}}{6 \text{ in } \underline{J}}$$

$$= -5.88 \approx -5.9$$

$$W = 5.9 \text{ J}$$

B

Q4 :

$$\Delta U + \Delta K = 0$$

$$8x^2 + 4x^4 = -\frac{1}{2}m(v_f^2 - v_i^2)$$

$$\text{at } x=1 \rightarrow v_f = 5$$

$$8 + 4 = -\frac{1}{2} \times 0.4 (25 - v_i^2)$$

$$12 = -0.2 (25 - v_i^2)$$

$$60 = v_i^2 - 25$$


$$v_i = 9.2$$

C

Q5: $U(x) = mgx + \frac{1}{2}kx^2$ $k \rightarrow \text{constant}$

$$\frac{-\partial U}{\partial x} = -(mg + kx) \quad \underline{E}$$

$$= -mg - kx$$

Q6: 

$h=5$ $P = \frac{W}{t} = \frac{-\Delta U}{t}$

$$\frac{-\Delta U}{t} = -d \cdot mg \sin 30 / t$$

$$= \frac{-5 \times 9.8 \times 600}{60} = 490$$

$d = \frac{h}{\sin \theta}$

$600 = \frac{10}{\sin 30} \times 60$

$P = 490$ A

Q7: none of the above E

Q8: When the block at high 50 m, its $\Delta U = \Delta U_{at 80} - \Delta U_{at 30}$

$$\Delta K = \Delta U$$

$$\Delta U_{at 80} = mgh = 6 \times 9.8 \times 80 = 4704$$

$$\Delta U_{at 30} = mgh = 6 \times 9.8 \times 30 = 1764$$

$$\Delta U_{at 50} = 4704 - 1764 = 2940 \quad \underline{C}$$

Q10: $F = \Delta P$

$\rightarrow +$ $P_{\text{before}} = P_{\text{after}}$

X-axis $mV \cos 30 = -mV \cos 30$


$$= -26$$

$\Delta P = P_f - P_i$

$$= -52 \hat{i} \quad \underline{D}$$

$P_i = 3 \times 10 \times \cos(30)$

$$= 15\sqrt{3} = 26$$



$V = 10$

at y-axi.

$$\Delta P = 0$$

$$+$$

$$\uparrow$$

$$V_i = V \sin 30 = 5$$

$$V_f = V \sin 30 = 5$$

$$\Delta P = m(V_f - V_i)$$

$$= 0$$

The University of Jordan / Department of Physics
First Semester 2015/2016
Physics 101/ Second Exam

Section number : KEY

Student name (بالعربية): _____

Lecturer name : _____

Student number : _____

✓ Some helpful information: gravitational acceleration $g = 9.8 \text{ m/s}^2$

Notes: Turn off your cell phone and put it out of sight. Keep your calculator on your own desk. Calculators cannot be shared. You have 75 minutes to complete your exam. Be sure to fill the box below with your final answers before the end of the exam.

	A	B	C	D	E		A	B	C	D	E
1						7					
2						8					
3						9					
4						10					
5						11					
6						12					

- A particle of mass (11 kg) is subject to two forces such that one force has a magnitude of 21 N directed east, and the other force has a magnitude of 39 N directed east-north, what is the magnitude of the particle's acceleration (in m/s^2)?
 (A) 2.8 **(B) 5.1** (C) 7.5 (D) 3.7 (E) 12
- An object of mass 4.0-kg is placed on top of an elevator floor. If the force exerted by the floor on the object is equal to 38 N. What is the acceleration of the elevator (in m/s^2)?
 (A) 0.8 upward (B) 0.8 downward (C) 1.3 upward (D) 1.3 downward **(E) 0.3 downward**
- A force of magnitude 20N directed in the positive x direction is acting on a particle and displacing it from the point (2m, -1m) to the point (4m, -3m). What is the work done by the force (in J)?
 (A) 60 **(B) 40** (C) 30 (D) 80 (E) 70
- A certain pendulum consists of a 1.5-kg mass swinging at the end of a string (length = 2.0 m). At the lowest point in the swing the tension in the string is equal to 20 N. To what maximum height (in cm) above this lowest point will the mass rise during its oscillation?
(A) 36 (B) 20 (C) 30 (D) 28 (E) 17
- A spring ($k = 600 \text{ N/m}$) is placed in a vertical position with its lower end supported by a horizontal surface. The upper end is compressed 20 cm, and a 4.0 kg block is placed on the compressed (مضغوط) spring. The system is then released from rest. How far above the point of release will the block rise (in cm)?
 (A) 20 **(B) 31** (C) 10 (D) 15 (E) 25

University of Jordan - Department of Physics - PHY 101
Second Exam - December 13, 2014 - (9:30 - 10:30) am

----- إسم الطالب: **KEY** ----- الرقم الجامعي: -----
----- إسم الدكتور: ----- رقم الشعبة أو وقت المحاضرة: -----

Given: ($g = 9.8 \text{ m/s}^2$)

Circle the letter of the correct answer ضغ دائرة حول حرف الإجابة الصحيحة

1.	A	B	C	D	E
2.	A	B	C	D	E
3.	A	B	C	D	E
4.	A	B	C	D	E
5.	A	B	C	D	E
6.	A	B	C	D	E
7.	A	B	C	D	E
8.	A	B	C	D	E
9.	A	B	C	XXX	XXX
10.	A	B	C	D	E
11.	A	B	C	D	E
12.	A	B	C	D	E

1.

A 1 kg particle undergoes a circular motion. At certain moment, the magnitude of the tangential and radial accelerations is 1.2 and 1.3 m/s² respectively. The magnitude of the total acceleration (in m/s²) for the particle at this moment is:

- A) 1.8 B) 1.2 C) 2.5 D) 0.1 E) 1.3

2.

A spring is stretched 5.00 cm from its equilibrium position. If this stretching requires 30.0 J of work, the spring constant (in kN/m) is:

- A) 24 B) 6 C) 12 D) 0.3 E) 1.3

3.

A 1.5 kg ball has a speed of 20 m/s when it is 15 m above the ground. The total energy (in J) of the ball is:

- A) 80 B) 300 C) 520 D) 220 E) 0

4.


A 1500 kg car accelerates from 0 to 25 m/s in 7 s. The average power delivered by the engine (1 hp = 746 W) is:

- A) 60 hp B) 80 hp C) 90 hp D) 70 hp E) 180 hp

5.

The coordinates of the center of mass for the system shown in Figure 1 are (L/4, -L/5). The coordinates of the 2-kg mass is:

- A) (-5L/8, 3L/10) B) (-11L/8, 9L/10)
C) (-5L/8, L/10) D) (-3L/8, L/10)
E) (-L/4, L/4)



6.

Consider a particle of mass m moving with linear momentum \vec{p} .

This particle is located at the vector position \vec{r} . The term $\left[\frac{d^2 \vec{r}}{dt^2} \times \frac{d\vec{p}}{dt} \right]$ gives:

- A) Force B) 0 C) Impulse D) Acceleration E) Velocity

7.

A 4 kg particle is subjected to a force acting in the x-direction, $F_x = (3+0.5x)$ N. The work (in J) done by the force as the particle moves from $x=0$ to $x=4$ m is:

- A) +20 B) -5 C) +16 D) 0 E) +5


8 & 9

8.

A 0.30 kg mass attached to the end of a string swings in a vertical circle ($R = 1.6$ m), as shown in **Figure 2**. At an instant when $\theta = 50^\circ$, the tension in the string is 8.0 N.

The magnitude of the resultant force (in N) on the mass at this instant is:

- A) 5.6 **B) 6.5** C) 6.1 D) 2.3 E) 5.1



(Figure 2)

9.

While the mass is passing the instant of the previous question ($\theta = 50^\circ$) and moving forward, the speed when $\theta = 51^\circ$ is:

A) Larger

B) The same

C) Smaller


10.

Figure 3 represents the magnitude of the net force (in N) exerted on a 3 kg mass. The magnitude of the impulse (in N.s) for the time interval between 2 and 5 s is:

- A) 0
D) 8

- B) 2** C) 1
E) 4

(Figure 3)



11.

A 10 kg object is dropped from rest. After falling a distance of 50 m, it has a speed of 26 m/s. The work (in kJ) done by the air resistive (friction) force on the object during this fall is:

A) -1.3

B) -1.5

C) -1.8

D) -2.0

E) -2.3

12.

A 0.28 kg ball has an elastic, head-on collision with a second ball that is initially at rest. The second ball moves off with half the original speed of the first ball. The mass (in kg) of the second ball is:

A) 0.14

B) 0.84

C) 0.42

D) 0.56

E) 0.28

Student's Name (In Arabic) _____


Registration #: _____ Sec# _____

Useful Information: **Some Results Are Rounded. CONSIDER (ACCELERATION DUE TO GRAVITY) $g = 9.8 \text{ m/s}^2$**

List your final answers in this table using **Capital Letters**.
Only the answer in this table will be graded

Question	Q1:	Q2:	Q3:	Q4:	Q5:	Q6:	Q7:	Q8:	Q9:	Q10:	Q11:	Q12:
Final Answer	D	B	A	B	C	D	E	C	A	C	B	D

Q1: A force F of 100 N is applied to a box of mass 5 kg moving on the floor as shown in the diagram. How much work (in J) is done by this force as the object moves 60 m?



- A) 4350 B) 7082
 C) 6241 D) 5196 E) 8042

Q2: If a spring is stretched 2.0 cm when a mass of 0.55 kg is hung vertically on it, then what is the force constant of the spring (in N/m)?

- A) 320 B) 270 C) 100 D) 84 E) 25

Q3: A box of mass 25.0 kg is placed on a rough horizontal surface. If the coefficient of kinetic friction between the surface and a box is 0.450, how much force is required to move the box at a constant speed across the surface (in N)?

- A) 110 B) 130 C) 140 D) 240 E) 270

Q4: If $F = (3\hat{i} - 8\hat{j}) \text{ N}$ is the only force acting on a 2.0-kg mass, what is the magnitude of the acceleration of the particle (m/s^2)?

- A) 1.5 B) 4.3 C) 6.5 D) 9.4 E) 7.2

Q5: An object is in a uniform circular motion. Which of the following statements must be true?
 حركة دائرية منتظمة


- A) The acceleration of the object is zero. B) The speed of the object is constant.
 C) The velocity of the object is constant. D) The acceleration of the object is constant.
 E) None of the above.

Answer Real

Q1	Q2	Q3	Q4	Q5	Q6	Q7	Q8	Q9	Q10
D	B	A	B	B	12	E	C	A	B
Q11	Q12	الاصحاب V.I.P							
B	D								

Q6: A ball attached to the end of a string of length of 4 m swings in a vertical circle as shown in the figure. The tension (in N) in the string when $\theta = 35^\circ$ is:

- A) 6.5
 B) 12
 C) 9.0
 D) 7.5
 E) 10




Q7: A particle moves along a circular path of radius 2.0 m. At an instant when the speed of the particle is equal to 3.0 m/s and changing at the rate of 5.0 m/s^2 , what is the magnitude of the total acceleration of the particle (in m/s^2)?

- A) 7.5
 B) 5.0
 C) 5.4
 D) 10
 E) 6.7

Q8: A spring under a compression x has a potential energy U_0 . When the compression is doubled $2x$, the potential energy stored in the spring is:

- A) U_0
 B) $2U_0$
 C) $4U_0$
 D) $6U_0$
 E) $8U_0$


Q9: A 2.0 kg block sits on a smooth (frictionless) plane inclined 20° above the horizontal. It is connected by a string to a 5.0 kg mass that hangs suspended from the other end of the string over a pulley as shown. Find the acceleration of the system.



- A) 6.0
 B) 8.0
 C) 10
 D) 12
 E) 15

Q10: The figure below represents the variation of the force (F) against the displacement (x) on a 5-kg mass. If the velocity at point B is 6 m/s, then the velocity (in m/s) at point A is:


- A) 2.1
 B) 5.1
 C) 7.8
 D) 8.15
 E) 9.2



Q11: An object of mass m rises vertically to a height h and then returns to its original point of projection. The work done by the gravitational force is:

- A) $-mgh$
 B) 0
 C) mgh
 D) $-2mgh$
 E) $2mgh$

Q12: A block starts from rest and slides down a frictionless inclined plane that is inclined 20° from the horizontal and is 2.53 meters long. What is the acceleration of the block (in m/s^2)?



- A) 10.2
 B) 9.40
 C) 7.32
 D) 3.35
 E) 2.53

Student's Name (In Arabic):..... **Key** Registration #:..... Sec#.....

Useful Information: **Some Results Are Rounded. CONSIDER (ACCELERATION DUE TO GRAVITY) $g = 9.8 \text{ m/s}^2$**

List your final answers in this table using **Capital Letters**.

Only the answer in this table will be graded

Question	Q1:	Q2:	Q3:	Q4:	Q5:	Q6:	Q7:	Q8:	Q9:	Q10:	Q11:	Q12:
Final Answer	D	C	E	D	B	B	B	A	D	C	A	B

Q1: An object is in a uniform circular motion. Which of the following statements must be true?


- A) The acceleration of the object is zero. B) The acceleration of the object is constant.
 C) The velocity of the object is constant. D) The speed of the object is constant.
 E) None of the above.

Q2: If $\mathbf{F} = (3\hat{i} - 8\hat{j}) \text{ N}$ is the only force acting on a 2.0-kg mass, what is the magnitude of the acceleration of the particle (m/s^2)?

- A) 1.5 B) 6.5 C) 4.3 D) 9.4 E) 7.2

Q3: A ball attached to the end of a string of length of 4 m swings in a vertical circle as shown in the figure. The tension (in N) in the string when $\theta = 35^\circ$ is:

- A) 6.5 B) 10 C) 9.0
 D) 7.5 E) 12



Q4: A particle moves along a circular path of radius 2.0 m. At an instant when the speed of the particle is equal to 3.0 m/s and changing at the rate of 5.0 m/s^2 , what is the magnitude of the total acceleration of the particle (m/s^2)?


- A) 7.5 B) 5.0 C) 5.4 D) 6.7 E) 4.5

Q5: A box of mass 25.0 kg is placed on a rough horizontal surface. If the coefficient of kinetic friction between the surface and a box is 0.450, how much force is required to move the box at a constant speed across the surface (in N)?

- A) 17.0 B) 110 C) 140 D) 240 E) 270


Q6: A block starts from rest and slides down a frictionless inclined plane that is inclined 20° from the horizontal and is 2.53 meters long. What is the acceleration of the block (in m/s^2)?

- A) 2.53 B) 3.35
C) 7.32 D) 9.40 E) 10.2




Q7: A force F of 100 N is applied to a box of mass 5 kg moving on the floor as shown in the diagram. How much work (in J) is done by this force as the object moves 60 m?

- A) 4350 B) 5196
C) 6241 D) 7082 E) 8042



Q8: The figure below represents the variation of the force (F) against the displacement (x) on a 5-kg mass. If the velocity at point B is 6 m/s, then the velocity (in m/s) at point A is:

- A) 5.1 B) 6.5
C) 7.8 D) 8.15 E) 9.2



Q9: A spring under a compression x has a potential energy U_0 . When the compression is doubled $2x$, the potential energy stored in the spring is:

- A) U_0 B) $2U_0$ C) $3U_0$ D) $4U_0$ E) $8U_0$


Q10: An object of mass m rises vertically to a height h and then returns to its original point of projection. The work done by the gravitational force is:

- A) mgh B) $-mgh$ C) 0 D) $-2mgh$ E) $2mgh$

Q11: If a spring is stretched 2.0 cm when a mass of 0.55 kg is hung vertically on it, then what is the force constant of the spring (in N/m)?

- A) 270 B) 320 C) 100 D) 84 E) 25

Q12: A 2.0 kg block sits on a smooth (frictionless) plane inclined 20° above the horizontal. It is connected by a string to a 5.0 kg mass that hangs suspended from the other end of the string over a pulley as shown. Find the acceleration of the system.



- A) 3.7 B) 6.0 C) 10 D) 12 E) 15



General Physics-1 (0302101) / Second Exam

Name (in Arabic): ----- **KEY ANSWER** ----- Instructor:-----

Registration No.: ----- Section: -----

- Choose the closest correct answer and fill the Answer Table. (Use $g = 9.8 \text{ m/s}^2$)

1. The tension in a string from which a 4.0-kg object is suspended in an elevator is equal to 44 N. What is the acceleration of the elevator?

- a. 11 m/s^2 upward. b. 1.2 m/s^2 downward. **c. 1.2 m/s^2 upward.**
d. 10 m/s^2 upward. e. 2.4 m/s^2 downward.

2. A book is placed on a chair. Then a videocassette is placed on the book. The floor exerts a normal force

- a. on all three. b. only on the book. **c. only on the chair.**
d. upwards on the chair and downwards on the book.
e. only on the objects that you have defined to be part of the system.

3. A 4.0-kg block slides down a 35° incline at a constant speed when a 16-N force is applied acting up and parallel to the incline. What is the coefficient of kinetic friction between the block and the surface of the incline?


- a. 0.20** b. 0.23 c. 0.26 d. 0.33 e. 0.41

4. A 4.0-kg mass on the end of a string rotates in a circular motion on a horizontal frictionless table. The mass has a constant speed of 2.0 m/s and the radius of the circle is 0.80 m. What is the magnitude of the resultant force acting on the mass?

- a. 40 N b. 30 N c. 44 N d. 0 N **e. 20 N**

5. A roller-coaster car has a mass of 500 kg when fully loaded with passengers. The car passes over a hill of radius 15 m, as shown. At the top of the hill, the car has a speed of 8.0 m/s. What is the force of the track on the car at the top of the hill?

- a. 7.0 kN up. b. 2.2 kN down. c. 2.8 kN down.
d. 2.8 kN up. e. 2.2 kN up.



6. A 30 kg child sitting 5.0 m from the center of a merry-go-round has a constant speed of 5.0 m/s. While she remains seated in the same spot and travels in a circle, the work the seat performs on her in one complete rotation is
a. zero. b. 150 J. c. 1500 J. d. 4700 J. e. 46,000 J.
7. A constant force of 15 N in the negative y direction acts on a particle as it moves from the origin to the point $(3\mathbf{i} + 3\mathbf{j} - 1\mathbf{k})$ m. How much work is done by the given force during this displacement?
 a. +30 J. **b. -45 J.** c. +45 J. d. -30 J. e. +75 J.
8. When a ball rises vertically to a height h and returns to its original point of projection, the work done by the gravitational force is
 a. $+2mgh$. b. $-mgh$. c. $+mgh$. d. $-2mgh$. **e. 0.**
9. The only force acting on a 2.0-kg body moving along the x axis is given by $F_x = (2x)$ N, where x is in m. If the velocity of the object at $x = 0$ is +3.0 m/s, how fast is it moving at $x = 2.0$ m?
 a. 5.0 m/s. **b. 3.6 m/s.** c. 4.1 m/s. d. 5.8 m/s. e. 2.8 m/s.
10. Carts A and B have equal masses and travel equal distances on straight frictionless tracks while a constant force F is applied to A, and a constant force $2F$ is applied to B. The relative amounts of work done by the two forces are related by
 a. $W_A = 4 W_B$. b. $W_A = 2 W_B$. c. $W_A = W_B$. **d. $W_B = 2 W_A$.** e. $W_B = 4 W_A$.

- Answer Table -

Fill the appropriate square of the correct answer with (X).

Q	a	b	c	d	e	Q	a	b	c	d	e
1						6					
2						7					
3						8					
4						9					
5						10					

University of Jordan
Faculty of Science
Department of Physics

Date: 31/12/2013
First Semester
Time: 4:00 – 5:00 pm

General Physics I – PHYS. 0302101
Makeup Second Exam

Name (In Arabic): **KEY ANSWER**
Student Number:

Instructor:
Section:

Constants: $g = 9.8 \text{ m/s}^2$

- Choose the closest correct answer and fill the Answer Table.

(Q1) A 0.5-kg mass attached to the end of a string swings in a vertical circle of radius equals 2.0 m. When the mass is at the lowest point on the circle, the speed of the mass is 12 m/s. The magnitude of the force (in N) of the string on the mass at this position is:

(A) 31 ; (B) 36 ; **(C) 41 ;** (D) 46 ; (E) 57 ;

(Q2) A particle moves in a circular path with constant speed. Its acceleration is:

(A) Zero ; (B) constantly increasing ; (C) constant in direction ;
(D) constant in magnitude and direction ; **(E) constant in magnitude ;**

(Q3) A 2.0-kg particle has an initial velocity of $(5\hat{i} - 4\hat{j}) \text{ m/s}$. Sometime later, its velocity is $(7\hat{i} + 3\hat{j}) \text{ m/s}$. How much work was done by the resultant force during this time interval, assuming no energy (in J) is lost in the process?

(A) 17 ; (B) 34 ; (C) 19 ; (D) 53 ; (E) 27 ;

(Q4) Equal amounts of work are performed on two bodies, A and B, initially at rest, and of masses M and 2M respectively. The relation between their speeds immediately after the work has been done on them is:

(A) $v_B = \sqrt{2}v_A$; (B) $v_B = 2 v_A$; (C) $v_A = v_B$; **(D) $v_A = \sqrt{2}v_B$;** (E) $v_A = 2 v_B$;

(Q5) A pendulum is made by letting a 2.0-kg object swing at the end of a string that has a length of 1.5 m. The maximum angle the string makes with the vertical as the pendulum swings is 30° . If air resistance is neglected, the speed (in m/s) of the object at the lowest point in its trajectory is:

(A) 1.6 ; **(B) 2.0 ;** (C) 2.5 ; (D) 2.7 ; (E) 3.1 ;

(Q6) A 10-N force acts on a 2.0-kg object initially at rest. The rate at which the force is doing work (in Watt) at time $t = 2.0 \text{ sec}$ is:

(A) 900 ; (B) 200 ; (C) 500 ; (D) 400 ; **(E) 100 ;**

(Q7) In a given displacement of a particle, its kinetic energy increases by 25 J while its potential energy decreases by 10 J. The work (in J) of the non-conservative forces acting on the particle during this displacement is:

(A) - 15 ; (B) +35 ; **(C) +15 ;** (D) - 35 ; (E) +55 ;

(Q8) A 3.0-kg ball with an initial velocity of $(4 \hat{i} + 3 \hat{j})$ m/s collides with a wall and rebounds with a velocity of $(-4 \hat{i} + 3 \hat{j})$ m/s. The impulse (in N.s) exerted on the ball by the wall is:

- (A) $-24 \hat{i}$; (B) $24 \hat{i}$; (C) $+18 \hat{j}$; (D) $-18 \hat{j}$; (E) $-16 \hat{i}$;

(Q9) A 2.0-kg object moving with a velocity of 5.0 m/s in the positive x direction collides with and sticks to an 8.0-kg object initially at rest. How much kinetic energy (in J) is lost in this collision?

- (A) 15; (B) 30; (C) 25; (D) 20; (E) 5;

(Q10) The turntable of a record player has an initial angular velocity of 8.0 rad/s at the moment when it is turned off. The turntable comes to rest 2.5 s after being turned off. Through how many radians does the turntable rotate after being turned off? Assume constant angular acceleration.

- (A) 12; (B) 8.0; (C) 10; (D) 16; (E) 6.8;

(Q11) Two points A and B are located on a disk that rotates about its axis. Point A is four times as far from the axis as point B . If the tangential speed of point B is equal to v , then the tangential speed of point A is:

- (A) v ; (B) $4v$; (C) $3v$; (D) $2v$; (E) $5v$;

(Q12) Two particles ($m_1 = 0.20$ kg, $m_2 = 0.30$ kg) are positioned at the ends of a 2.0- m long rod of negligible mass. The moment of inertia (in $kg.m^2$) of this system about an axis perpendicular to the rod and through the center of mass is:

- (A) 0.38; (B) 0.75; (C) 1.2; (D) 0.48; (E) 1.7;

-Answer Table-

Fill the appropriate square of the correct answer with (X).

Q's	A	B	C	D	E	Q's	A	B	C	D	E
1						7					
2						8					
3						9					
4						10					
5						11					
6						12					

General Physics I – PHYS. 0302101
Second Exam

Name (In Arabic): **KEY ANSWER**
Student Number:

Instructor:
Section:

Constants: $g = 9.8 \text{ m/s}^2$

* Choose the closest correct answer and fill the Answer Table.

(Q1) An airplane moves 100 m/s as it travels around a vertical circular loop which has a 1.0-km radius. The magnitude of the resultant force (in kN) on the 70-kg pilot of this plane at the bottom of this loop is:

(A) 0.70 ; (B) 1.37 ; (C) 2.1 ; (D) 1.3 ; (E) 1.58 ;

(Q2) An object (a) of mass m flies in a horizontal circle of radius R at a speed v . Another object (b) has the same mass m and flies in a horizontal circle of radius R at a speed of $v/2$. Then the ratio of the centripetal acceleration of the object (a) to that of object (b) is:

(A) 0.25 ; (B) 0.5 ; (C) 1.0 ; (D) 2.0 ; (E) 4.0 ;

(Q3) Single conservative force acting on an object moving along the x axis is given by: $F_x = (14x - 3x^2) \text{ N}$, where x is in m . The Change in potential energy ΔU (in J) done by this force as the object moves from $x = -1 \text{ m}$ to $x = +2 \text{ m}$ is :

(A) - 20.1 ; (B) + 38.0 ; (C) - 12.0 ; (D) + 16.0 ; (E) - 28.0 ;

(Q4) A 12-kg block on a horizontal frictionless surface is attached to a light spring (force constant = 700 N/m). The block is initially at rest at its equilibrium position when a force of magnitude 80 N acting parallel to the surface is applied to the block. The speed (in m/s) of the block when it is 13 cm from its equilibrium position is:

(A) 0.55 ; (B) 0.68 ; (C) 0.78 ; (D) 0.86 ; (E) 0.90 ;

(Q5) A constant force of 10 N in the negative y direction acts on a particle as it moves from the origin to the point $(3\hat{i} + 3\hat{j} - 1\hat{k}) \text{ m}$. The work (in J) done by the given force during this displacement is:

(A) - 45 ; (B) - 30 ; (C) - 60 ; (D) +30 ; (E) +12 ;

(Q6) A 2.0-kg block slides down a plane (inclined at 40° with the horizontal) at a constant speed of 5.0 m/s. The Power (in W) at which the gravitational force doing on the block is:

(A) zero; (B) - 55.2; (C) + 78.7; (D) + 94.5; (E) + 63.0;

(Q7) Three particles are placed in the xy plane. A 30 g particle is located at $(3, 4) m$, a 40 g particle is located at $(-2, -2) m$. Where a 20 g particle must be placed (in m) so that the center of mass of the three-particle system is at the Origin?

- (A) $(-0.5, -2.0)$; (B) $(1, 0)$; (C) $(2.5, 2)$; (D) $(-3, -14)$; (E) $(0, -2)$;

(Q8) A 2.0-kg object is moving along the x -axis. Its speed increases from 30 m/s to 40 m/s during a 5.0-s time interval. The magnitude of the average total force (in N) acting on the object during this time interval is:

- (A) 2.0; (B) 3.0; (C) 4.0; (D) 5.0; (E) 6.0;

(Q9) A ball falls to the ground from height H and bounces to height h . Momentum is conserved in the ball-earth system

- (A) only if $h > H$; (B) only if $h = 0$; (C) only if $h = H$;
 (D) only if $h \leq H$; (E) only if $h \geq H$;

(Q10) At $t = 0$, a wheel rotating about a fixed axis at a constant angular acceleration has an angular velocity of 2.0 rad/s . Two seconds later it has turned through 5.0 complete revolutions. The angular acceleration (in rad/s^2) of this wheel is:

- (A) 15.7; (B) 13.7; (C) 9.7; (D) 7.7; (E) 5.7;

(Q11) A wheel rotating about a fixed axis has an angular position given by $\theta = 3 - 2t^3$, where θ is measured in radians and t in seconds. The angular velocity (in rad/s) of the wheel at $t = 2.0$ s is:

- (A) -24 ; (B) -38 ; (C) -54 ; (D) -62 ; (E) -96 ;

Q12) A disk with a radius of $2.0 m$ whose moment of inertia is $50 \text{ kg}\cdot\text{m}^2$ rotates uniformly by angular acceleration of 6.0 rad/s^2 . The net force (in N) acting tangent to the circumference of this disk is:

- (A) 75; (B) 100; (C) 115; (D) 135; (E) 150;

Answer Table

Fill the appropriate square of the correct answer with (X).

Q's	A	B	C	D	E	Q's	A	B	C	D	E
1						7					
2						8					
3						9					
4						10					
5						11					
6						12					

General Physics I – PHYS. 0302101
Makeup Second Exam

Name (In Arabic): **KEY ANSWER**
Student Number:

Instructor:
Section:

Constants: $g = 9.8 \text{ m/s}^2$

- Choose the closest correct answer and fill the Answer Table.

(Q1) A 0.5-kg mass attached to the end of a string swings in a vertical circle of radius equals 2.0 m. When the mass is at the lowest point on the circle, the speed of the mass is 12 m/s. The magnitude of the force (in N) of the string on the mass at this position is:

- (A) 31 ; (B) 36 ; (C) 41 ; (D) 46 ; (E) 57 ;

(Q2) A particle moves in a circular path with constant speed. Its acceleration is:

- (A) Zero ; (B) constantly increasing ; (C) constant in direction ;
(D) constant in magnitude and direction ; (E) constant in magnitude ;

(Q3) A 2.0-kg particle has an initial velocity of $(5\hat{i} - 4\hat{j}) \text{ m/s}$. Sometime later, its velocity is $(7\hat{i} + 3\hat{j}) \text{ m/s}$. How much work was done by the resultant force during this time interval, assuming no energy (in J) is lost in the process?

- (A) 17 ; (B) 34 ; (C) 19 ; (D) 53 ; (E) 27 ;

(Q4) Equal amounts of work are performed on two bodies, A and B, initially at rest, and of masses M and 2M respectively. The relation between their speeds immediately after the work has been done on them is:

- (A) $v_B = \sqrt{2}v_A$; (B) $v_B = 2v_A$; (C) $v_A = v_B$; (D) $v_A = \sqrt{2}v_B$; (E) $v_A = 2v_B$;

(Q5) A pendulum is made by letting a 2.0-kg object swing at the end of a string that has a length of 1.5 m. The maximum angle the string makes with the vertical as the pendulum swings is 30° . If air resistance is neglected, the speed (in m/s) of the object at the lowest point in its trajectory is:

- (A) 1.6 ; (B) 2.0 ; (C) 2.5 ; (D) 2.7 ; (E) 3.1 ;

(Q6) A 10-N force acts on a 2.0-kg object initially at rest. The rate at which the force is doing work (in Watt) at time $t = 2.0 \text{ sec}$ is:

- (A) 900 ; (B) 200 ; (C) 500 ; (D) 400 ; (E) 100 ;

(Q7) In a given displacement of a particle, its kinetic energy increases by 25 J while its potential energy decreases by 10 J. The work (in J) of the non-conservative forces acting on the particle during this displacement is:

- (A) - 15 ; (B) +35 ; (C) +15 ; (D) - 35 ; (E) +55 ;

(Q8) A 3.0-kg ball with an initial velocity of $(4 \hat{i} + 3 \hat{j})$ m/s collides with a wall and rebounds with a velocity of $(-4 \hat{i} + 3 \hat{j})$ m/s. The impulse (in N.s) exerted on the ball by the wall is:

- (A) $-24 \hat{i}$; (B) $24 \hat{i}$; (C) $+18 \hat{j}$; (D) $-18 \hat{j}$; (E) $-16 \hat{i}$;

(Q9) A 2.0-kg object moving with a velocity of 5.0 m/s in the positive x direction collides with and sticks to an 8.0-kg object initially at rest. How much kinetic energy (in J) is lost in this collision?

- (A) 15; (B) 30; (C) 25; (D) 20; (E) 5;

(Q10) The turntable of a record player has an initial angular velocity of 8.0 rad/s at the moment when it is turned off. The turntable comes to rest 2.5 s after being turned off. Through how many radians does the turntable rotate after being turned off? Assume constant angular acceleration.

- (A) 12; (B) 8.0; (C) 10; (D) 16; (E) 6.8;

(Q11) Two points A and B are located on a disk that rotates about its axis. Point A is four times as far from the axis as point B . If the tangential speed of point B is equal to v , then the tangential speed of point A is:

- (A) v ; (B) $4v$; (C) $3v$; (D) $2v$; (E) $5v$;

(Q12) Two particles ($m_1 = 0.20$ kg, $m_2 = 0.30$ kg) are positioned at the ends of a 2.0-m long rod of negligible mass. The moment of inertia (in $kg.m^2$) of this system about an axis perpendicular to the rod and through the center of mass is:

- (A) 0.38; (B) 0.75; (C) 1.2; (D) 0.48; (E) 1.7;

-Answer Table-

Fill the appropriate square of the correct answer with (X).

Q's	A	B	C	D	E	Q's	A	B	C	D	E
1						7					
2						8					
3						9					
4						10					
5						11					
6						12					



Physics 1 Second Exam



Zenon
مركز التعليم الإلكتروني

Mech Family

Civilium
مركز التعليم الإلكتروني

Madar
ask, believe and realize

22

Q1: A car ($m = 2234 \text{ kg}$) and its velocity is 25 m/s collide with a tree, it needs 0.26 seconds to stopped them;

Find force applied from tree on car ..

Q2: An object start moving in a circular path from rest, after (10.9) sec., the object has a rate with (12000 rev/min) ,

Find its angular acceleration ..

Q3: An object moves in a circular path, the angular displacement is given with $(3 - 2t^3)$;

Find angular acceleration after 1 sec.

Q4: A ball ($m = 22 \text{ kg}$) is thrown to a wall with velocity (31 m/s) and it rebounds after (0.11 sec) with velocity (18 m/s) then

Find the force applied

Q5: An object is moving from rest and after 10 sec. it has an angular velocity (50 rad/s) , if its moment of Inertia

$(I = 9 \text{ kg.m}^2)$... Find its Torque ...


Q6: If an object is moving in constant velocity, on a circular path which radius $(r = 0.01 \text{ m})$, and its angular velocity (5 rev./sec.) then ... Find acceleration ..

Q7: Find I (I for one object $= \frac{ML^2}{3}$) ...



Q8 :- If an object (7 kg) is moving with velocity $(8\hat{i} + 9\hat{j})$
 Find its momentum (magnitude)


Q9 :- If an object is moving as shown (Tension = 40 N) ...
 Find angular velocity ??



Q10 :- An object is moving with angular velocity (20 rad/s)
 and after (5 sec.) the angular velocity (12 rad/s) ...
 Find angular displacement ... ?

Q11 :- Two object is moving with velocities ($m_1 = 3243$,
 $m_2 = 1300$)
 $(v_1 = 49 \text{ m/s}, v_2 = -6.5 \text{ m/s})$...
 They collide and start moving as one object ...
 Find their velocity after collision ... ??

Q12 :- Find I about O ...



** Solu. 1 g - $\Delta P = F_{avg} \cdot \Delta t$
 $(mv_f - mv_i)$
 $\Rightarrow 2234(0) - 2234(25) = F_{avg} \cdot 0.26$

$F_{avg} = 2.1 \times 10^5$

** Solu. 2 g -

$1200 \frac{\text{rev.}}{\text{min.}} \cdot \frac{1 \text{ min.}}{60} \cdot \frac{2\pi}{1 \text{ rev.}} = 1256.6 \text{ rad/s.}$

$\omega_f = \omega_i + \alpha t$

$\Rightarrow 1256.6 = 0 + 10.5(\alpha) \Rightarrow \alpha = 120 \text{ rad/s}^2$

** Solu. 3 g -

$\theta = 3 - 2t^3$

$\omega = -6t^2 // \alpha = -12t \Rightarrow \alpha = -12 \text{ rad/s}^2$
 $t=1$

** Solu. 4 g -

$\Delta P = F_{avg} \cdot \Delta t$

$mv_f - mv_i = F_{avg} \cdot \Delta t$

$\begin{matrix} -18 \text{ m/s} \\ 31 \text{ m/s} \end{matrix} \Bigg| \Delta t = 0.11 \text{ s.}$

$22(-18 - 31) = F \cdot 0.11 \Rightarrow F = -9800 \text{ N}$

** Solu. 5 g -

$\tau = I\alpha$

$= 9 \cdot 5$

$= 45 \frac{\text{kg} \cdot \text{m}^2}{\text{s}^2}$

$\omega_t = \omega_i + t\alpha$

$50 = 0 + 10\alpha$

$\alpha = 5 \text{ rad/s}^2$

** Solu. 6 g -

$v = r\omega$

$v = 0.01 \cdot 31.4$

$= 0.314 \text{ m/sec.}$

$a_t = \frac{v^2}{r} = \frac{0.314^2}{0.01} = 9.86 \text{ m/s}^2$

** Solu. 7 g -

$I = I_1 + I_2$

$= \frac{M_1 L^2}{3} + \frac{M_2 L^2}{3} = \frac{2^2}{3} (0.8 + 0.3)$

$I = 1.47 \text{ kg} \cdot \text{m}^2$

$$\text{** Sol. 8 } \vec{a} = m\vec{V}$$

$$= 7(8\hat{i} - 19\hat{j}) = 56\hat{i} + 63\hat{j}$$

$$P = \sqrt{56^2 + 63^2} = 84.3 \frac{\text{kg} \cdot \text{m}}{\text{s}^2}$$

** Solu. 9

$$F = \frac{Mv^2}{r}$$

$$40 = \frac{2 \cdot v^2}{5}$$

$$r = 10 \text{ m/s}$$

$$v = r \cdot \omega$$

$$\omega = \frac{10}{5} = 2 \text{ rad/sec.}$$

** Solu. 10

$$\omega_f = \omega_i + \alpha t$$

$$\alpha = -1.6 \text{ rad/s}^2$$

$$\Delta\phi = \omega_i t + \frac{1}{2} \alpha t^2$$

$$= 20 \cdot 5 - 0.5 \cdot 1.6 \cdot 25$$

$$= 80 \text{ rad.}$$

** Solu. 11

$$P_i = P_f$$

$$M_1 v_1 + M_2 v_2 = (M_1 + M_2) v_2$$

$$3243 \cdot 49 - 1300 \cdot 65 = (3243 + 1300) v_2$$

$$v_2 = 16.4 \text{ m/s}$$

** Solu. 12

$$I = \sum M r^2$$

$$= 5^2 (7 + 6 + 6 + 10) = 725 \text{ kg} \cdot \text{m}^2$$

Good Luck



GENERAL PHYSICS 1:(0302101)
SECOND EXAM

NOTE: Acceleration due to gravity, $g=9.8 \text{ m/s}^2$

الاسم باللغة العربية: _____ الرقم الجامعي: _____

اسم المدرس: _____ الشعبة: _____ رقم الجلوس: _____

- 1) A 15kg block is placed on a rough horizontal surface of $\mu_s = 0.3$. The block is kept in equilibrium as shown in the figure.

The maximum hanging mass for which the system will remain in equilibrium is:

- (a) 2.6 (b) 25.5 (c) 42.1 (d) 76.4 (e) 4.3



- 2) A conical pendulum is formed by attaching a small ball to a 1.2m string. The ball swings with uniform velocity around a horizontal circle of radius 30cm as shown in the figure. The velocity (m.s^{-1}) of the ball is:

- (a) 11.5 (b) 0.72 (c) 0.87 (d) 3.4 (e) 0.52



- 3) A 4kg particle experiences a net force along the x-axis given by $F=3x^2 - 6$, where F is in Newton and x is in meters. If the particle starts to move from rest at $x=0$, the power (w) delivered to the particle when it is at $x=4\text{m}$ is:


- (a) 168 (b) 150 (c) 476 (d) 345 (e) 188

- 4) A force $F=(5y^2 \text{ N.m}^{-2})\hat{j}$ is applied to a particle. The work done (J) by the force on the particle as it moves along a straight line from (2, 3) to (5, 5) is:


- (a) 527 (b) 20 (c) 1466 (d) 163 (e) 200

- 5) This graph represents the power developed by a motor. The energy (J) expended by the motor in time interval $t=10\text{s}$ to $t=30\text{s}$ is:


- (a) 200 (b) 100 (c) 0.5 (d) 600 (e) 500




- 6) A 2.2 kg block placed on a frictionless 20° inclined plane. A force of 16 N acting parallel to the incline as shown in the figure. The acceleration (m/s^2) of the block is:
 (a) 2.0 down the incline (b) 5.3 up the incline (c) 2.0 up the incline
 (d) 3.9 down the incline (e) 3.9 up the incline




- 7) An object attached to the end of a string swings in a vertical circle (يتحرك في دائرة عمودية) of radius 1.2 m, as shown in the figure. At an instant when $\theta = 30^\circ$, the speed of the object is 6.0 m/s and the tension in the string is 38 N. The mass (kg) of the object is:
 (a) 2.0 (b) 1.5 (c) 1.8 (d) 1.3 (e) 0.80




- 8) A block of mass 5.0 kg is moving with 3.0 m/s on a rough horizontal surface (coefficient of kinetic friction = 0.40) when it collides with a spring, as shown in the figure. The spring is compressed a maximum distance of 0.20 m. The spring constant (N/m) is:
 (a) 1020 (b) 1804 (c) 2196 (d) 361 (e) 929



- 9) A 1.2-kg mass is projected down a rough circular track (radius = 2.0 m) as shown below. The speed of the mass at point A is 3.5 m/s, and at point B, it is 6.0 m/s. How much work is done on the mass between A and B by the force of friction?
 (a) -9.3 J (b) -7.3 J (c) -8.1 J (d) -10.8 J (e) -24 J



- 10) A 4 kg mass is placed on a rough horizontal surface. Two forces in the same plane act on the mass as shown in the figure. The magnitude of the force $F(N)$ that enable (تمكنه من) the 4kg mass to accelerate with $(3m.s^{-2})\hat{j}$ is:
 (a) 13.4 (b) 7.5 (c) 6.7 (d) 4.8 (e) 10.0



Answer Table

Q.No.	A	B	C	D	e
1					
2					
3					
4					
5					
6					
7					
8					
9					
10					

Notes: Turn off your cell phone and put it out of sight. Keep your calculator on your own desk. Calculators cannot be shared. You have 75 minutes to complete your exam.

Be sure to fill the box below with your final answers before the end of the exam.

	A	B	C	D	E	A	B	C	D	E
1		/					/			
2							/			
3		/					/			
4		/					/			
5		/					/			
6							/			

- A particle of mass (1 kg) is subject to two forces such that one force has a magnitude of 21 N directed east, and the other force has a magnitude of 39 N directed east-north. What is the magnitude of the particle's acceleration (in m/s^2)?

(A) 2.8 (B) 5.1 (C) 7.5 (D) 3.7 (E) 12
- An object of mass 4.0-kg is placed on top of an elevator floor. If the force exerted by the floor on the object is equal to 38 N. What is the acceleration of the elevator (in m/s^2)?

(A) 0.8 upward (B) 0.8 downward (C) 1.3 upward (D) 1.3 downward (E) 0.3 downward
- A force of magnitude 20N directed in the positive x direction is acting on a particle and displacing it from the point (2m, -1m) to the point (4m, -5m). What is the work done by the force (in J)?

(A) 60 (B) 40 (C) 30 (D) 80 (E) 70
- A certain pendulum consists of a 1.5-kg mass swinging at the end of a string (length = 2.0 m). At the lowest point in the swing the tension in the string is equal to 20 N. To what maximum height (in cm) above this lowest point will the mass rise during its oscillation?

(A) 36 (B) 20 (C) 30 (D) 28 (E) 17
- A spring ($K = 600 \text{ N/m}$) is placed in a vertical position with its lower end supported by a horizontal surface. The upper end is compressed 20 cm, and a 4.0 kg block is placed on the compressed (مضغوط) spring. The system is then released from rest. How far above the point of release will the block rise (in cm)?

(A) 20 (B) 31 (C) 10 (D) 15 (E) 25

6. A potential energy function for a two-dimensional force is of the form $U = 3x^2y$. Find the force that acts at the point (1, 1).

(A) $\vec{F} = -12\hat{i} - 3\hat{j}$ (B) $\vec{F} = -6\hat{j}$ (C) $\vec{F} = -24\hat{i} - 12\hat{j}$ (D) $\vec{F} = 6\hat{i} - 3\hat{j}$ (E) $\vec{F} = -6\hat{i}$

7. A 6.0-kg block slides along a horizontal surface. If $\mu_s = 0.20$ for the block and surface, at what rate is the friction force doing work on the block (in W) at an instant when its speed is 4.0 m/s ?

(A) -63 (B) -47 (C) -50 (D) +25 (E) -55

8. A particle of mass (1.5 kg) is moving on the xy -axis with an acceleration given as $a = (6.0\hat{i} + 5.0\hat{j}) \text{ m/s}^2$. What is the speed of the particle in (m/s) at the moment it reaches $x = 4.0 \text{ m}$, given that the particle started motion from origin with initial velocity 2.0 m/s ?

(A) 10.1 (B) 14.7 (C) 11.8 (D) 13.1 (E) 9.5


9. An airplane moves at constant speed of 140 m/s as it travels around a vertical circular loop which has a 1.0-km radius. What is the magnitude of the net force causing the centripetal acceleration on the 71-kg pilot (in N)?

(A) 1000 (B) 1392 (C) 1200 (D) 1310 (E) 1022

10. A roller-coaster car has a mass of 400 kg when fully loaded with passengers ($\sim 5j$). At the bottom of a circular dip of radius 40 m (as shown in the figure) the car has a speed of 16 m/s . What is the magnitude of the force the track exerts on the car at the bottom of the dip (in kN)?

(A) 10.1 (B) 9.7 (C) 8.1 (D) 13.1 (E) 6.5

11. What is the magnitude of the tension in the string (in N) if $M = 2.0 \text{ kg}$ in the figure shown? Assume the surface is frictionless.



(A) 21 (B) 19.7 (C) 32.2 (D) 42.9 (E) 56.5

12. A box of mass (42 kg) is placed on top of a rough horizontal surface whose coefficients of friction are ($\mu_s = 0.5, \mu_k = 0.4$). If a man tried to push the box by applying a force of (210 N), what would be the magnitude of the friction force (in N)?


(A) 210 (B) 247 (C) 220 (D) 155 (E) 230

List your final answers in this table using Capital Letters
Only the answer in this table will be graded

Question	Q1:	Q2:	Q3:	Q4:	Q5:	Q6:	Q7:	Q8:	Q9:	Q10:
Final Answer	R	A	D	B	A	D	R	R	D	C

Q1: For the vectors shown in the figure, express vector \vec{S} in terms of vectors \vec{M} and \vec{N} .

- A) $\vec{M} = \vec{S} - \vec{N}$
- B) $\vec{S} = \vec{M} - \vec{N}$
- C) $\vec{M} = \vec{N} - \vec{S}$
- D) $\vec{M} + \vec{S} + \vec{N} = 0$
- E) $\vec{N} = \vec{S} + \vec{M}$



Q2: An airplane undergoes the following displacements: First, it flies 66 km in a direction 30° east of north. Next, it flies 49 km due south. Finally, it flies 100 km 30° north of west. How far (in km) the airplane ends up from its starting point.

- A) 79
- B) 81
- C) 73
- D) 86
- E) 93

Q3: What is the angle between the vector $\vec{A} = +3\hat{i} - 2\hat{j} - 3\hat{k}$ and the $+\hat{y}$ -axis?


- A) 25°
- B) 65°
- C) 90°
- D) 115°
- E) 155°

Q4: The figure shows the position of an object as a function of time. What is the average velocity (in m/s) of the object during the time interval from time $t = 3.0 \text{ s}$ and time $t = 9.0 \text{ s}$.

- A) 0.33
- B) 0.33
- C) 0.66
- D) 0.66
- E) 0.99



$$\Delta v = \frac{\Delta x}{\Delta t} = \frac{1-3}{9-3} = \frac{-2}{6} = -\frac{1}{3}$$



Q5: The velocity of an object as a function of time is given by $v(t) = 2.00 \text{ m/s} + (3.00 \text{ m/s})t - (1.0 \text{ m/s}^2)t^2$. Determine the acceleration (in m/s^2) of the object at time $t = 5.00 \text{ s}$.

- A) -7.00 B) -2.00 C) 0.00 D) 2.00 E) 7.00

Q6: A car starts from rest and accelerates with a constant acceleration of 1.00 m/s^2 for $t = 3.00 \text{ s}$. The car then continues for 5.00 s at constant velocity. How far (in m) has the car traveled from its starting point?

- A) 4.50 B) 7.00 C) 15.0 D) 19.5 E) 25.0

Q7: A ball is thrown upward at time $t = 0.00 \text{ s}$, from a point on a roof 70 m above the ground and experiences negligible air resistance. The ball rises, then falls and strikes the ground. The initial velocity of the ball is 28.5 m/s . The velocity of the ball (in m/s) when it is 39 m above the ground is:

- A) -45 B) -38 C) 38 D) 45 E) -23

Q8: An object has a position given by $\vec{r} = [2.0 \text{ m} + (5.00 \text{ m/s})\hat{j}] + [3.0 \text{ m} - (2.00 \text{ m/s}^2)t^2]\hat{i}$, where quantities are in SI units. What is the speed (in m/s) of the object at time $t = 2.00 \text{ s}$?

- A) 13.0 B) 9.43 C) 7.58 D) 6.40 E) 1.42

Q9: A child throws a ball with an initial speed of 8.00 m/s at an angle of 40.0° above the horizontal. The ball leaves her hand 1.00 m above the ground and experience negligible air resistance. How far (in m) from where the child is standing does the ball hit the ground?

- A) 1.67 B) 3.80 C) 5.05 D) 6.39 E) 7.46

Q10: A ball is tied to the end of a cable of negligible mass. The ball is rotated in a circle with a radius 2.00 m making 7.00 revolutions every 10.0 seconds. What is the magnitude of the acceleration of the ball (in m/s^2)?

- A) 74.2 B) 67.9 C) 38.7 D) 29.3 E) 14.8